



Fermion Hierarchies from String Theory

Lucas Leung

based on arXiv:2410.17704 and arXiv:2507.XXXXX

In collaboration with: Andrei Constantin, Kit Fraser-Taliente, Thomas Harvey, Andre Lukas and Luca Nutricati

International School of Subnuclear Physics 2025 Erice - 20th June 2025

Fermion Hierarchies from String Theory

Lucas Leung

based on arXiv:2410.17704 and arXiv:2507.XXXXX



In collaboration with: Andrei Constantin, Kit Fraser-Taliente, Thomas Harvey, Andre Lukas and Luca Nutricati

International School of Subnuclear Physics 2025 Erice - 20th June 2025



Fermion Masses and Mixings

Why do we have fermion mass hierarchies?



$$V_{\text{CKM}} \sim \begin{pmatrix} 1 & 0.2 & 0.2^3 \\ 0.2 & 1 & 0.2^2 \\ 0.2^3 & 0.2^2 & 1 \end{pmatrix}$$

stolen from Admir Greljo's La Thuile and Matthew McCullough's Erice talk

Froggatt-Nielsen Mechanism

- **Froggatt and Nielsen** [1979] proposed using horizontal symmetries $U(1)_H$ to explain flavour structures

$$y_{ij} = a_{ij} \langle \phi \rangle^{n_{ij}}$$

O(1) coefficient

$\leftarrow U(1)_H$ neutral

Q_1	Q_2	Q_3	$\bar{d}_1$	$\bar{d}_2$	$\bar{d}_3$	$\bar{u}_1$	$\bar{u}_2$	$\bar{u}_3$
(3)	(2)	(0)	(3)	(2)	(2)	(3)	(1)	(0)

+

ϕ
(-1)

+
 Set $\langle \phi \rangle \sim \lambda$

Froggatt-Nielsen Mechanism

- **Froggatt and Nielsen** [1979] proposed using horizontal symmetries $U(1)_H$ to explain flavour structures

$$y_{ij} = a_{ij} \langle \phi \rangle^{n_{ij}} \quad \begin{array}{l} \text{O(1) coefficient} \\ \text{U(1)}_H \text{ neutral} \end{array}$$

Q_1	Q_2	Q_3	$\bar{d}_1$	$\bar{d}_2$	$\bar{d}_3$	$\bar{u}_1$	$\bar{u}_2$	$\bar{u}_3$
(3)	(2)	(0)	(3)	(2)	(2)	(3)	(1)	(0)

+

ϕ
(-1)

+
Set $\langle \phi \rangle \sim \lambda$

$$M^d \sim \langle H_d \rangle \begin{pmatrix} \lambda^6 & \lambda^5 & \lambda^5 \\ \lambda^5 & \lambda^4 & \lambda^4 \\ \lambda^3 & \lambda^2 & \lambda^2 \end{pmatrix}, \quad M^u \sim \langle H_u \rangle \begin{pmatrix} \lambda^6 & \lambda^4 & \lambda^3 \\ \lambda^5 & \lambda^3 & \lambda^2 \\ \lambda^3 & \lambda & 1 \end{pmatrix}.$$

Froggatt-Nielsen Mechanism

- **Froggatt and Nielsen** [1979] proposed using horizontal symmetries $U(1)_H$ to explain flavour structures

$$\phi^6 Q_1 H_d \bar{d}_1$$

$$y_{ij} = a_{ij} \langle \phi \rangle^{n_{ij}}$$

O(1) coefficient

$\leftarrow U(1)_H$ neutral

Q_1	Q_2	Q_3	$\bar{d}_1$	$\bar{d}_2$	$\bar{d}_3$	$\bar{u}_1$	$\bar{u}_2$	$\bar{u}_3$
(3)	(2)	(0)	(3)	(2)	(2)	(3)	(1)	(0)

+


ϕ
(-1)

+
Set $\langle \phi \rangle \sim \lambda$

$$M^d \sim \langle H_d \rangle \begin{pmatrix} \lambda^6 & \lambda^5 & \lambda^5 \\ \lambda^5 & \lambda^4 & \lambda^4 \\ \lambda^3 & \lambda^2 & \lambda^2 \end{pmatrix}, \quad M^u \sim \langle H_u \rangle \begin{pmatrix} \lambda^6 & \lambda^4 & \lambda^3 \\ \lambda^5 & \lambda^3 & \lambda^2 \\ \lambda^3 & \lambda & 1 \end{pmatrix}.$$



String Model Building

- most well-studied QG theory
- maybe best theory to understand HEP



String Model Building

- most well-studied QG theory
- maybe best theory to understand HEP





Want to focus on heterotic
 $E_8 \times E_8$ on smooth CYs

String Model Building

- most well-studied QG theory
- maybe best theory to understand HEP

Why Strings?

Intuition to how different 'nuts and bolts' in a HE theory 'interact' to give Standard Model + answer other pheno questions.



Want to focus on heterotic $E_8 \times E_8$ on smooth CYs

U(1) symmetries from string theory

- Low-energy effective theories have gauge symmetry

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times \mathcal{G}$$

$$\text{where } \mathcal{G} \cong U(1)^n / \mathbb{Z}n$$

field	SM rep	name	SU(5)	$\mathcal{G}$ charge pattern	SU(5) $\times$ $\mathcal{G}$
Q	$(\mathbf{3}, \mathbf{2})_1$	LH quark	$\mathbf{10}$	$\mathbf{e}_a$	$\mathbf{10}_a$
u	$(\bar{\mathbf{3}}, \mathbf{1})_{-4}$	RH u -quark			
e	$(\mathbf{1}, \mathbf{1})_6$	RH electron			
d	$(\bar{\mathbf{3}}, \mathbf{1})_2$	RH d -quark	$\bar{\mathbf{5}}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}$
L	$(\mathbf{1}, \mathbf{2})_{-3}$	LH lepton			
H^d	$(\mathbf{1}, \mathbf{2})_{-3}$	down-Higgs	$\bar{\mathbf{5}}^{H^d}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}^{H^d}$
H^u	$(\mathbf{1}, \mathbf{2})_3$	up-Higgs	$\mathbf{5}^{H^u}$	$-\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{5}_{a,b}^{H^u}$
ϕ	$(\mathbf{1}, \mathbf{1})_0$	pert. FN scalar	$\mathbf{1}$	$\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{1}_{a,b}$
Φ	$(\mathbf{1}, \mathbf{1})_0$	non-pert. FN scalar	$\mathbf{1}$	$\mathbf{k} = (k_1, \dots, k_f)$	$\mathbf{1}$

U(1) symmetries from string theory

- Low-energy effective theories have gauge symmetry

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times \mathcal{G}$$

$$\text{where } \mathcal{G} \cong U(1)^n / \mathbb{Z}n$$

field	SM rep	name	SU(5)	$\mathcal{G}$ charge pattern	SU(5) $\times$ $\mathcal{G}$
Q	$(\mathbf{3}, \mathbf{2})_1$	LH quark	$\mathbf{10}$	$\mathbf{e}_a$	$\mathbf{10}_a$
u	$(\bar{\mathbf{3}}, \mathbf{1})_{-4}$	RH u -quark			
e	$(\mathbf{1}, \mathbf{1})_6$	RH electron			
d	$(\bar{\mathbf{3}}, \mathbf{1})_2$	RH d -quark	$\bar{\mathbf{5}}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}$
L	$(\mathbf{1}, \mathbf{2})_{-3}$	LH lepton			
H^d	$(\mathbf{1}, \mathbf{2})_{-3}$	down-Higgs	$\bar{\mathbf{5}}^{H^d}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}^{H^d}$
H^u	$(\mathbf{1}, \mathbf{2})_3$	up-Higgs	$\mathbf{5}^{H^u}$	$-\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{5}_{a,b}^{H^u}$
ϕ	$(\mathbf{1}, \mathbf{1})_0$	pert. FN scalar	$\mathbf{1}$	$\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{1}_{a,b}$
Φ	$(\mathbf{1}, \mathbf{1})_0$	non-pert. FN scalar	$\mathbf{1}$	$\mathbf{k} = (k_1, \dots, k_f)$	$\mathbf{1}$

**Different to traditional
Froggatt-Nielsen!**

U(1) symmetries from string theory

- Low-energy effective theories have gauge symmetry

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times \mathcal{G}$$

where $\mathcal{G} \cong U(1)^n / \mathbb{Z}n$

1 - Discrete Quotients

field	SM rep	name	SU(5)	$\mathcal{G}$ charge pattern	SU(5) $\times$ $\mathcal{G}$
Q	$(\mathbf{3}, \mathbf{2})_1$	LH quark	$\mathbf{10}$	$\mathbf{e}_a$	$\mathbf{10}_a$
u	$(\bar{\mathbf{3}}, \mathbf{1})_{-4}$	RH u -quark			
e	$(\mathbf{1}, \mathbf{1})_6$	RH electron			
d	$(\bar{\mathbf{3}}, \mathbf{1})_2$	RH d -quark	$\bar{\mathbf{5}}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}$
L	$(\mathbf{1}, \mathbf{2})_{-3}$	LH lepton			
H^d	$(\mathbf{1}, \mathbf{2})_{-3}$	down-Higgs	$\bar{\mathbf{5}}^{H^d}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}^{H^d}$
H^u	$(\mathbf{1}, \mathbf{2})_3$	up-Higgs	$\mathbf{5}^{H^u}$	$-\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{5}_{a,b}^{H^u}$
ϕ	$(\mathbf{1}, \mathbf{1})_0$	pert. FN scalar	$\mathbf{1}$	$\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{1}_{a,b}$
Φ	$(\mathbf{1}, \mathbf{1})_0$	non-pert. FN scalar	$\mathbf{1}$	$\mathbf{k} = (k_1, \dots, k_f)$	$\mathbf{1}$

Different to traditional Froggatt-Nielsen!

U(1) symmetries from string theory

- Low-energy effective theories have gauge symmetry

$$SU(3)_c \times SU(2)_L \times U(1)_Y \times \mathcal{G}$$

where $\mathcal{G} \cong U(1)^n / \mathbb{Z}n$

1 - Discrete Quotients

field	SM rep	name	SU(5)	$\mathcal{G}$ charge pattern	SU(5) $\times$ $\mathcal{G}$
Q	$(\mathbf{3}, \mathbf{2})_1$	LH quark	$\mathbf{10}$	$\mathbf{e}_a$	$\mathbf{10}_a$
u	$(\bar{\mathbf{3}}, \mathbf{1})_{-4}$	RH u -quark			
e	$(\mathbf{1}, \mathbf{1})_6$	RH electron			
d	$(\bar{\mathbf{3}}, \mathbf{1})_2$	RH d -quark	$\bar{\mathbf{5}}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}$
L	$(\mathbf{1}, \mathbf{2})_{-3}$	LH lepton			
H^d	$(\mathbf{1}, \mathbf{2})_{-3}$	down-Higgs	$\bar{\mathbf{5}}^{H^d}$	$\mathbf{e}_a + \mathbf{e}_b$	$\bar{\mathbf{5}}_{a,b}^{H^d}$
H^u	$(\mathbf{1}, \mathbf{2})_3$	up-Higgs	$\mathbf{5}^{H^u}$	$-\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{5}_{a,b}^{H^u}$
ϕ	$(\mathbf{1}, \mathbf{1})_0$	pert. FN scalar	$\mathbf{1}$	$\mathbf{e}_a - \mathbf{e}_b$	$\mathbf{1}_{a,b}$
Φ	$(\mathbf{1}, \mathbf{1})_0$	non-pert. FN scalar	$\mathbf{1}$	$\mathbf{k} = (k_1, \dots, k_f)$	$\mathbf{1}$

Different to traditional Froggatt-Nielsen!


2 - Specific Charges for scalar + SM fields

Goal

- Want to understand **fermion hierarchies** using **U(1)s from string theory**

Top-down Approach

- Use symmetry patterns of known string models - compute mass hierarchies
- **Difficult - satisfy other phenomenological properties (Higgs? Neutrinos?)!**




String Theory

Fermion Hierarchies

Bottom-up Approach

- Use charge patterns from string theory as constraints of effective theory
- **Large number of possible patterns - analyse using Machine Learning & Genetic Algorithms**



Examples

Top-down Approach

Downstairs Spectrum

<i>Matter fields</i>	$\mathbf{10}_1, \mathbf{10}_2, \mathbf{10}_5, \bar{\mathbf{5}}_{1,3}, \bar{\mathbf{5}}_{2,3}, \bar{\mathbf{5}}_{4,5}$
<i>Higgs fields</i>	$H_{3,5}, \bar{H}_{3,5}$
<i>Moduli</i>	$\Phi_1, \dots, \Phi_5$
	$\phi_{1,2}, \phi_{2,1}, \phi_{1,4}, \phi_{5,1}, \phi_{2,3},$ $\phi_{2,5}, \phi_{3,4}, \phi_{4,3}, \phi_{3,5}, \phi_{5,3}, \phi_{5,4}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{3,5} \phi_{5,1}^2 & \phi_{2,1} \Phi_2 & \phi_{3,5} \phi_{5,1} \\ \phi_{2,1} \Phi_2 & \Phi_2 & \phi_{2,5} \Phi_2 \\ \phi_{3,5} \phi_{5,1} & \phi_{2,5} \Phi_2 & \phi_{3,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{2,1} \phi_{4,3} & \phi_{4,3} & \phi_{2,5} \\ \phi_{4,3} & 0 & 0 \\ \phi_{2,5} \phi_{4,3} & 0 & \Phi_1 \end{pmatrix},$$

Bottom-up Approach

Downstairs Spectrum

<i>Matter fields</i>	$\mathbf{10}_1, \mathbf{10}_2, \mathbf{10}_5, 3 \bar{\mathbf{5}}_{3,4}$
<i>Higgs fields</i>	$H_{4,5}, \bar{H}_{4,5}$
<i>Moduli</i>	Φ_1, Φ_2
	$\phi_{4,5}, \phi_{5,1}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{4,5} \phi_{5,1}^2 & \phi_{4,5}^2 \phi_{5,1}^2 \Phi_1 & \phi_{4,5} \phi_{5,1} \\ \phi_{4,5}^2 \phi_{5,1}^2 \Phi_1 & \phi_{4,5}^3 \phi_{5,1}^2 \Phi_1^2 & \phi_{4,5}^2 \phi_{5,1} \Phi_1 \\ \phi_{4,5} \phi_{5,1} & \phi_{4,5}^2 \phi_{5,1} \Phi_1 & \phi_{4,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 \\ \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 \\ \Phi_2 & \Phi_2 & \Phi_2 \end{pmatrix}$$

Examples

1 - Models exist!

Top-down Approach

Downstairs Spectrum

<i>Matter fields</i>	$\mathbf{10}_1, \mathbf{10}_2, \mathbf{10}_5, \bar{\mathbf{5}}_{1,3}, \bar{\mathbf{5}}_{2,3}, \bar{\mathbf{5}}_{4,5}$
<i>Higgs fields</i>	$H_{3,5}, \bar{H}_{3,5}$
<i>Moduli</i>	$\Phi_1, \dots, \Phi_5$
	$\phi_{1,2}, \phi_{2,1}, \phi_{1,4}, \phi_{5,1}, \phi_{2,3},$ $\phi_{2,5}, \phi_{3,4}, \phi_{4,3}, \phi_{3,5}, \phi_{5,3}, \phi_{5,4}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{3,5} \phi_{5,1}^2 & \phi_{2,1} \Phi_2 & \phi_{3,5} \phi_{5,1} \\ \phi_{2,1} \Phi_2 & \Phi_2 & \phi_{2,5} \Phi_2 \\ \phi_{3,5} \phi_{5,1} & \phi_{2,5} \Phi_2 & \phi_{3,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{2,1} \phi_{4,3} & \phi_{4,3} & \phi_{2,5} \\ \phi_{4,3} & 0 & 0 \\ \phi_{2,5} \phi_{4,3} & 0 & \Phi_1 \end{pmatrix},$$

Bottom-up Approach

Downstairs Spectrum

<i>Matter fields</i>	$\mathbf{10}_1, \mathbf{10}_2, \mathbf{10}_5, 3 \bar{\mathbf{5}}_{3,4}$
<i>Higgs fields</i>	$H_{4,5}, \bar{H}_{4,5}$
<i>Moduli</i>	Φ_1, Φ_2
	$\phi_{4,5}, \phi_{5,1}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{4,5} \phi_{5,1}^2 & \phi_{4,5} \phi_{5,1}^2 \Phi_1 & \phi_{4,5} \phi_{5,1} \\ \phi_{4,5}^2 \phi_{5,1}^2 \Phi_1 & \phi_{4,5}^3 \phi_{5,1}^2 \Phi_1^2 & \phi_{4,5}^2 \phi_{5,1} \Phi_1 \\ \phi_{4,5} \phi_{5,1} & \phi_{4,5}^2 \phi_{5,1} \Phi_1 & \phi_{4,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 \\ \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 \\ \Phi_2 & \Phi_2 & \Phi_2 \end{pmatrix}$$

Examples

1 - Models exist!

Top-down Approach

Downstairs Spectrum

Matter fields	$10_1, 10_2, 10_5, \bar{5}_{1,3}, \bar{5}_{2,3}, \bar{5}_{4,5}$
Higgs fields	$H_{3,5}, \bar{H}_{3,5}$
Moduli	$\Phi_1, \dots, \Phi_5$
	$\phi_{1,2}, \phi_{2,1}, \phi_{1,4}, \phi_{5,1}, \phi_{2,3},$ $\phi_{2,5}, \phi_{3,4}, \phi_{4,3}, \phi_{3,5}, \phi_{5,3}, \phi_{5,4}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{3,5} \phi_{5,1}^2 & \phi_{2,1} \Phi_2 & \phi_{3,5} \phi_{5,1} \\ \phi_{2,1} \Phi_2 & \Phi_2 & \phi_{2,5} \Phi_2 \\ \phi_{3,5} \phi_{5,1} & \phi_{2,5} \Phi_2 & \phi_{3,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{2,1} \phi_{4,3} & \phi_{4,3} & \phi_{2,5} \\ \phi_{4,3} & 0 & 0 \\ \phi_{2,5} \phi_{4,3} & 0 & \Phi_1 \end{pmatrix},$$

Bottom-up Approach

Downstairs Spectrum

Matter fields	$10_1, 10_2, 10_5, 3 \bar{5}_{3,4}$
Higgs fields	$H_{4,5}, \bar{H}_{4,5}$
Moduli	Φ_1, Φ_2
	$\phi_{4,5}, \phi_{5,1}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{4,5} \phi_{5,1}^2 & \phi_{4,5} \phi_{5,1}^2 \Phi_1 & \phi_{4,5} \phi_{5,1} \\ \phi_{4,5}^2 \phi_{5,1}^2 \Phi_1 & \phi_{4,5}^3 \phi_{5,1}^2 \Phi_1^2 & \phi_{4,5}^2 \phi_{5,1} \Phi_1 \\ \phi_{4,5} \phi_{5,1} & \phi_{4,5}^2 \phi_{5,1} \Phi_1 & \phi_{4,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 \\ \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 \\ \Phi_2 & \Phi_2 & \Phi_2 \end{pmatrix}$$

2 - a lot of extra fields!

Examples

1 - Models exist!

Top-down Approach

Downstairs Spectrum

Matter fields	$10_1, 10_2, 10_5, \bar{5}_{1,3}, \bar{5}_{2,3}, \bar{5}_{4,5}$
Higgs fields	$H_{3,5}, \bar{H}_{3,5}$
Moduli	$\Phi_1, \dots, \Phi_5$
	$\phi_{1,2}, \phi_{2,1}, \phi_{1,4}, \phi_{5,1}, \phi_{2,3}, \phi_{2,5}, \phi_{3,4}, \phi_{4,3}, \phi_{3,5}, \phi_{5,3}, \phi_{5,4}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{3,5} \phi_{5,1}^2 & \phi_{2,1} \Phi_2 & \phi_{3,5} \phi_{5,1} \\ \phi_{2,1} \Phi_2 & \Phi_2 & \phi_{2,5} \Phi_2 \\ \phi_{3,5} \phi_{5,1} & \phi_{2,5} \Phi_2 & \phi_{3,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{2,1} \phi_{4,3} & \phi_{4,3} & \phi_{2,5} \\ \phi_{4,3} & 0 & 0 \\ \phi_{2,5} \phi_{4,3} & 0 & \Phi_1 \end{pmatrix},$$

Bottom-up Approach

3 - charge constraints!

Downstairs Spectrum

Matter fields	$10_1, 10_2, 10_5, 3 \bar{5}_{3,4}$
Higgs fields	$H_{4,5}, \bar{H}_{4,5}$
Moduli	Φ_1, Φ_2
	$\phi_{4,5}, \phi_{5,1}$

Yukawa Insertions

$$\Lambda^u \sim \begin{pmatrix} \phi_{4,5} \phi_{5,1}^2 & \phi_{4,5} \phi_{5,1}^2 \Phi_1 & \phi_{4,5} \phi_{5,1} \\ \phi_{4,5}^2 \phi_{5,1}^2 \Phi_1 & \phi_{4,5}^3 \phi_{5,1}^2 \Phi_1^2 & \phi_{4,5}^2 \phi_{5,1} \Phi_1 \\ \phi_{4,5} \phi_{5,1} & \phi_{4,5}^2 \phi_{5,1} \Phi_1 & \phi_{4,5} \end{pmatrix}$$

$$\Lambda^d \sim \begin{pmatrix} \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 & \phi_{5,1} \Phi_2 \\ \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 & \phi_{4,5} \phi_{5,1} \Phi_1 \Phi_2 \\ \Phi_2 & \Phi_2 & \Phi_2 \end{pmatrix}$$

2 - a lot of extra fields!

Conclusions & Outlook

- Can construct **viable models** with **fermionic hierarchies** using **U(1) symmetries** from heterotic string theory!

Conclusions & Outlook

- Can construct **viable models** with **fermionic hierarchies** using **U(1) symmetries** from heterotic string theory!

Top-down Approach

- String models with **good** fermionic hierarchies found!!!
- Need to satisfy other pheno properties (correct Higgs scale $\sqrt{}$, neutrinos?)

Bottom-up Approach

- Constructed GA environment to obtain **list of viable models!**
- Guidance to top-down model building (topological constraints)!

Conclusions & Outlook

- Can construct **viable models** with **fermionic hierarchies** using **U(1) symmetries** from heterotic string theory!

Top-down Approach

- String models with **good** fermionic hierarchies found!!!
- Need to satisfy other pheno properties (correct Higgs scale $\sqrt{}$, neutrinos?)

Link?
↔


Bottom-up Approach

- Constructed GA environment to obtain **list of viable models!**
- Guidance to top-down model building (topological constraints)!

Backup Slides

Genetic Algorithms


- A family of optimisation-search algorithms.
- Two parts: **Environment** + **Evolution**



Evolution

Environment


“0010101100”	$m_u = 0.002$	→	0
“0000100100”	$m_u = 0.03$	→	-1.18
“1110010000”	$m_u = 150$	→	-4.88
Genotype	Phenotype		Fitness



Bottom-up Results - Scans (Perturbative Only)

n	Fixed charges	N_ϕ	Env Size	States visited	Full scan	Models	Inequiv. spectra
(1, 1, 1, 1, 1)	$10_1, 10_2, 10_5, \bar{5}_{4,5}^H$	2	10^9		Yes	0	0
		3	10^{11}	10^8		0	0
		4	10^{13}	10^8		301	4
		5	10^{16}	10^9		29213	289
(1, 1, 1, 2)	$10_1, 10_2, 10_4, \bar{5}_{4,4}^H$	2	10^8		Yes	0	0
		3	10^{10}		Yes	98	1
		4	10^{12}	10^8		18825	55
		5	10^{14}	10^8		320557	449
(1, 1, 1, 2)	$10_2, 10_3, 10_4, \bar{5}_{1,4}^H$	2	10^8		Yes	0	0
		3	10^{10}		Yes	56	1
		4	10^{12}	10^8		11538	128
		5	10^{14}	10^8		259175	1128
(1, 1, 1, 2)	$10_1, 10_2, 10_3, \bar{5}_{1,4}^H$	2	10^8		Yes	0	0
		3	10^{10}		Yes	70	3
		4	10^{12}	10^8		8110	63
		5	10^{14}	10^8		204148	500
(1, 1, 1, 2)	$10_1, 10_3, 10_4, \bar{5}_{1,2}^H$	2	10^8		Yes	0	0
		3	10^{10}		Yes	0	0
		4	10^{12}	10^8		0	0
		5	10^{14}	10^8		0	0
(1, 1, 3)	$10_1, 10_2, 10_3, \bar{5}_{3,3}^H$	2	10^6		Yes	8	1
		3	10^8		Yes	1218	18
		4	10^{10}		Yes	22734	81
		5	10^{12}	10^8		154532	234
(1, 2, 2)	$10_1, 10_2, 10_3, \bar{5}_{3,3}^H$	2	10^6		Yes	0	0
		3	10^8		Yes	0	0
		4	10^{10}		Yes	0	0
		5	10^{12}	10^8		0	0

Total number of inequiv. spectra obtained against states visited



Environment size
 $\sim \mathcal{O}(10^{14})$

Bottom-up Example

Spectrum
 $\mathbf{10}_1, \mathbf{10}_2, \mathbf{10}_5; 3\bar{\mathbf{5}}_{1,2}; H_{4,5}^u, H_{4,5}^d$
 $\phi_{5,1}, \phi_{3,5}, \phi_{1,2}, \phi_{4,1}, \phi_{4,5}$

Optimise coefficients - $\epsilon = 0.554$ and

$$\begin{pmatrix} 1.090 & 2.282 & 1.896 \\ 0.961 & 2.027 & 1.979 \\ 1.966 & 2.978 & 2.648 \end{pmatrix}$$


up-quark sector

$$\begin{pmatrix} 1.379 & 1.843 & 0.947 \\ 2.708 & 1.726 & 2.063 \\ 1.530 & 2.526 & 0.680 \end{pmatrix}$$

down-quark sector

$$\begin{pmatrix} 1.064 & 2.051 & 1.707 \\ 1.183 & 2.628 & 2.262 \\ 0.514 & 1.623 & 0.706 \end{pmatrix}$$

lepton sector



Yukawa Textures

up sector:

$$\begin{pmatrix} \phi_{5,1}\phi_{4,1} & \phi_{5,1}\phi_{1,2}\phi_{4,1} & \phi_{4,1} \\ \phi_{5,1}\phi_{1,2}\phi_{4,1} & \phi_{5,1}\phi_{1,2}^2\phi_{4,1} & \phi_{1,2}\phi_{4,1} \\ \phi_{4,1} & \phi_{1,2}\phi_{4,1} & \phi_{4,5} \end{pmatrix}$$

down sector:

$$\begin{pmatrix} \phi_{5,1}\phi_{3,5} & \phi_{5,1}\phi_{3,5} & \phi_{5,1}\phi_{3,5} \\ \phi_{5,1}\phi_{3,5}\phi_{1,2} & \phi_{5,1}\phi_{3,5}\phi_{1,2} & \phi_{5,1}\phi_{3,5}\phi_{1,2} \\ \phi_{3,5} & \phi_{3,5} & \phi_{3,5} \end{pmatrix}$$

Compute Quantities

Higgs VEV	$\langle H \rangle = 174 \text{ GeV}$		
Quark	$m_u \text{ (MeV)}$	$m_c \text{ (GeV)}$	$m_t \text{ (GeV)}$
Mass	2.16	1.27	173
Quark	$m_d \text{ (MeV)}$	$m_s \text{ (MeV)}$	$m_b \text{ (GeV)}$
Mass	4.70	93.9	4.18
Lepton	$m_e \text{ (MeV)}$	$m_\mu \text{ (MeV)}$	$m_\tau \text{ (GeV)}$
Mass	0.511	106	1.78

$$|V_{\text{CKM}}| \simeq \begin{pmatrix} 0.970 & 0.242 & 0.00359 \\ 0.242 & 0.969 & 0.0447 \\ 0.00733 & 0.0443 & 0.999 \end{pmatrix}$$